



HHS Public Access

Author manuscript

Nature. Author manuscript; available in PMC 2013 January 19.

Published in final edited form as:

Nature. 2012 July 19; 487(7407): 370–374. doi:10.1038/nature11184.

Proto-genes and *de novo* gene birth

Anne-Ruxandra Carvunis^{1,2}, Thomas Rolland¹, Ilan Wapinski³, Michael A. Calderwood¹, Muhammed A. Yildirim⁴, Nicolas Simonis^{1,†}, Benoit Charlotiaux^{1,5}, César A. Hidalgo⁶, Justin Barbette¹, Balaji Santhanam¹, Gloria A. Brar⁷, Jonathan S. Weissman⁷, Aviv Regev^{8,9}, Nicolas Thierry-Mieg², Michael E. Cusick¹, and Marc Vidal^{1,¶}

¹Center for Cancer Systems Biology (CCSB) and Department of Cancer Biology, Dana-Farber Cancer Institute, and Department of Genetics, Harvard Medical School, Boston, MA 02215, USA.

²UJF-Grenoble 1 / CNRS / TIMC-IMAG UMR 5525, Computational and Mathematical Biology Group, Grenoble, F-38041, France.

³Department of Systems Biology, Harvard Medical School, Boston, MA 02115, USA.

⁴Center for International Development and Harvard University, Cambridge, MA 02138, USA.

⁵Unit of Animal Genomics, GIGA-R and Faculty of Veterinary Medicine, University of Liege, 4000 Liege, Wallonia-Brussels Federation, Belgium.

⁶The MIT Media Lab, Massachusetts Institute of Technology, Cambridge, MA 02142, USA.

⁷Howard Hughes Medical Institute, Department of Cellular and Molecular Pharmacology, University of California, San Francisco, and California Institute for Quantitative Biosciences, San Francisco, CA 94158, USA.

⁸Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA.

⁹Howard Hughes Medical Institute, Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02139, USA.

Abstract

Novel protein-coding genes can arise either through re-organization of pre-existing genes or *de novo*^{1,2}. Processes involving re-organization of pre-existing genes, notably following gene

Users may view, print, copy, download and text and data- mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use: http://www.nature.com/authors/editorial_policies/license.html#terms

Author Information Correspondence and requests for materials should be addressed to M.V. (marc_vidal@dfci.harvard.edu).

[¶]Correspondence and requests for materials should be addressed to marc_vidal@dfci.harvard.edu.

[†]Present address: Laboratoire de Bioinformatique des Génomes et des Réseaux (BiGRé), Campus Plaine, Free University of Brussels, 1050 Brussels, Wallonia-Brussels Federation, Belgium.

Full methods and any associated references are available in the online version of the paper at www.nature.com/nature.

The authors declare no competing financial interests.

Author Contributions A.-R.C., I.W., M.E.C. and M.V. conceived the project. A.-R.C. led the project and performed most of the analyses. T.R. evaluated cross-species transfer events, optimized the ribosome footprint analysis pipeline and assisted in other analyses. I.W. designed the conservation level tool and calculated most of the purifying selection statistics. M.A.Y. aligned the sequencing reads. B.S. predicted disordered and transmembrane regions and assisted in the cross-species transfer analyses. N.S. and B.C. assisted in analyses. G.A.B. and J.S.W. shared their expertise in ribosome footprinting data analysis and provided the meiosis ribosome footprinting raw and processed data. A.-R.C., T.R., M.E.C. and M.V. designed the figures. All authors contributed to writing the manuscript.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

duplication, have been extensively described^{1,2}. In contrast, *de novo* gene birth remains poorly understood, mainly because translation of sequences devoid of genes, or “non-genic” sequences, is expected to produce insignificant polypeptides rather than proteins with specific biological functions^{1,3-6}. Here, we formalize an evolutionary model according to which functional genes evolve *de novo* through transitory proto-genes⁴ generated by widespread translational activity in non-genic sequences. Testing this model at genome-scale in *Saccharomyces cerevisiae*, we detect translation of hundreds of short species-specific open reading frames (ORFs) located in non-genic sequences. These translation events appear to provide adaptive potential⁷, as suggested by their differential regulation upon stress and by signatures of retention by natural selection. In line with our model, we establish that *S. cerevisiae* ORFs can be placed within an evolutionary continuum ranging from non-genic sequences to genes. We identify ~1,900 candidate proto-genes among *S. cerevisiae* ORFs and find that *de novo* gene birth from such a reservoir may be more prevalent than sporadic gene duplication. Our work illustrates that evolution exploits seemingly dispensable sequences to generate adaptive functional innovation.

Both genome-wide surveys and analyses of individual cases have shown that *de novo* gene birth has occurred throughout the evolution of many lineages, potentially impacting species-specific adaptations and evolutionary radiations^{1,2,5,6,8,9}. Genes are thought to emerge *de novo* when non-genic sequences become transcribed, acquire ORFs and the corresponding non-genic transcripts access the translation machinery^{1,2,4,5,8}. However, it is hard to reconcile this proposed mechanism with expectations that non-genic sequences should lack translational activity and, even if translated, should encode insignificant polypeptides^{1,3,4,6}. Evidence of associations between non-genic >transcripts and ribosomes has suggested that non-genic sequences may occasionally be translated, which could provide raw material for natural selection⁶. It has also been speculated that genes that originate *de novo* could initially be simple and gradually become more complex over evolutionary time⁴. These ideas are consistent with reports showing that genes that emerged recently are shorter, less expressed and more rapidly diverging than other genes^{1,10-13}. We developed an integrative evolutionary model whereby *de novo* gene birth proceeds through intermediate and reversible proto-gene stages, mirroring the well-described pseudo-gene stages of gene death (Fig. 1a)¹⁴.

We investigated this model at genome-scale in the context of *de novo* gene birth in *Saccharomyces cerevisiae*^{8,10}. In *S. cerevisiae*, a minimal length threshold of 300 nucleotides was originally used to delineate ORFs likely to be genes from non-genic ORFs occurring by chance in non-genic sequences¹⁵. The resulting gene catalogue has undergone numerous adjustments¹⁶, with currently ~6,000 ORFs annotated as genes and ~261,000 unannotated ORFs containing at least three codons considered non-genic ORFs (Supplementary Fig. 1). Nongenic sequences are broadly transcribed in *S. cerevisiae*¹⁷, their overexpression is mostly non-toxic¹⁸, and the corresponding transcripts can associate with ribosomes, often at AUGs^{6,19}. We reasoned that translation of non-genic ORFs could be more common than expected. Such translation events would not systematically lead to *de novo* gene birth, since the corresponding polypeptides would not necessarily have specific biological functions. Instead, upon translation, non-genic ORFs would become proto-genes (Fig. 1b). Proto-genes would provide adaptive potential⁶ by exposing genetic variations that

are usually hidden in non-genic sequences. A subset of proto-genes could occasionally be retained over evolutionary time, for instance if providing an advantage to the organism under specific environmental conditions. Retained proto-genes could gradually evolve the characteristics of genes, while other proto-genes might lose the ability to be translated. Such a reservoir of proto-genes would allow evolutionary innovations to be attempted without affecting existing genes.

This evolutionary model leads to the following predictions: i) the structural and functional characteristics of *S. cerevisiae* ORFs (e.g. length, expression level or sequence composition) should reflect an evolutionary continuum ranging from non-genic ORFs to genes; ii) many non-genic ORFs should be translated; iii) ORFs that emerged recently should occasionally have adaptive functions retained by natural selection.

To examine these predictions, we estimated the order of emergence of *S. cerevisiae* ORFs (Fig. 1c). Annotated ORFs were classified into 10 groups based on their conservation throughout the Ascomycota phylogeny (Supplementary Fig. 2). Of ~6,000 annotated ORFs, ~2% are found only in *S. cerevisiae* (ORFs₁) (Supplementary Fig. 2)¹⁰ and ~12% are found only in the four closely related *Saccharomyces sensu stricto* species (ORFs₁₋₄). The ~88% of annotated ORFs found outside of this group (ORFs₅₋₁₀) are well characterized and can confidently be considered genes. ORFs₁₋₄ are poorly characterized and their annotation as genes is debatable (Supplementary Fig. 2)^{16,20}. The weak conservation of ORFs₁₋₄ suggests that they emerged recently, which we corroborated using gene duplication events to control for relative time of emergence (Supplementary Fig. 3). We estimate that over 97% of ORFs₁₋₄ originated *de novo* rather than by cross-species transfer, which could also explain their weak conservation (Supplementary Information). ORFs₁₋₄ often partially overlap ORFs₅₋₁₀, which seems incompatible with cross-species transfer, or preferentially lie within subtelomeric regions whose instability may facilitate *de novo* emergence (Supplementary Fig. 4). In addition to classifying ORFs₁₋₁₀, we assigned a conservation level of 0 to ~108,000 unannotated ORFs longer than 30 nucleotides and free from overlap with annotated features on the same strand (ORFs₀) (Supplementary Information). ORFs₀ and ORFs₁₋₄ constituted our initial list of candidate proto-genes.

To test the evolutionary continuum prediction, we first verified that ORF conservation level correlates positively with length and expression level (Fig. 2a and Supplementary Fig. 5)^{1,10-12}. These correlations suggest that genes evolve from non-genic ORFs that lengthen and increase in expression level over evolutionary time. A negative correlation between ORF length and expression level²¹ was observed among ORFs₅₋₁₀, but not among ORFs₁₋₄ (Supplementary Fig. 5). Thus, some ORFs may increase in expression level at different rates than they increase in length over evolutionary time. Lengthening of ORFs could occur by loss of stop codons, possibly following translational read-through, by shift of start codons or by duplication followed by fusion with other ORFs^{10,22}. Increase in ORF expression level could be mediated by recruitment of existing regulatory elements¹. The proportion of ORFs located in the vicinity of transcription factor binding sites increases with conservation level, suggesting that novel regulatory elements could also emerge (Fig. 2a)¹.

In line with a study of codon evolution in metazoans²³, we observed a positive correlation between codon usage bias and conservation level (Fig. 2b). Relative abundances of amino acids in proteins encoded by ORFs₁₋₄ show levels intermediate between those in proteins encoded by ORFs₅₋₁₀ and in hypothetical translation products of ORFs₀ (Fig. 2c), similar to observations in bacteria²⁴. Likely due to this biased sequence composition, ORFs₁₋₄ exhibit a higher hydrophobicity, a higher tendency to form transmembrane regions and a lower propensity for intrinsic structural disorder¹⁰ than ORFs₅₋₁₀ (Fig. 2d). Taken together, our observations support the existence of an evolutionary continuum ranging from non-genic ORFs to genes.

To assess the extent of non-genic translation, we searched for signatures of translation of ORFs₀ at genome-scale in a ribosome footprinting dataset generated in both rich and starvation conditions²⁵. In this dataset, ~1% of sequencing reads could not be mapped to ORFs₁₋₁₀. We developed a stringent pipeline to detect unequivocal translation signatures for ORFs₀ located on transcripts associated with ribosomes (Fig. 3a and Supplementary Fig. 6). We found that 1,139 of ~108,000 ORFs₀ show such evidence of translation (ORFs₀⁺). This number is significantly higher than expected if the ribosome footprinting assay was non-specific, or if the presence of ribosomes on non-genic transcripts was unrelated to the presence of ORFs₀ (Fig. 3b). These ORFs₀⁺ are enriched in adenine at position -3 from the start codon, which likely favours translation initiation (Fig. 3c and Supplementary Information). We verified that ORFs₀⁺ did not originate from gene duplication or cross-species transfer and are not genes that have failed to be annotated due to their short length (Supplementary Information). The 1,139 ORFs₀⁺ therefore appear to be translated non-genic ORFs.

We detected strong differential translation of ORFs₀⁺ and ORFs₁₋₄ in starvation or rich conditions, whereas most ORFs₅₋₁₀ are translated in both conditions (Fig. 3d and Supplementary Fig. 6). We found that the binding sites of four transcription factors involved in mating and stress response are preferentially located close to ORFs₀⁺ and ORFs₁₋₄ (Supplementary Table 1) and that ORFs₁₋₄ are enriched in the Gene Ontology term “response to stress” (Supplementary Table 2). Recently emerged ORFs may provide adaptive functions in response to environmental stress.

Retention by natural selection was measured by comparing the genome sequences of eight *S. cerevisiae* strains to evaluate the tendency of ORF sequences to be purged of non-synonymous mutations (purifying selection) relative to expectations under neutral evolution. Most ORFs₀⁺ and ORFs₁₋₄ do not exhibit a significant deviation from neutral evolution, yet ~3% of ORFs₀⁺ and 9-25% of ORFs₁₋₄ appear under purifying selection (Fig. 3e). This fraction increases with conservation level, in line with the proposed evolutionary continuum (Supplementary Fig. 7 and Supplementary Information). Our observations suggest that recently emerged ORFs occasionally acquire adaptive functions that are retained by natural selection, in agreement with findings in primates and with evolutionary models derived from inter-species comparisons^{12,13,26}.

Overall, our results show that *de novo* gene birth could proceed through proto-genes. From the initial comprehensive set of candidate proto-genes (all ORFs₀ and ORFs₁₋₄), we

excluded ORFs₀ that appear to lack translation signatures according to our stringent pipeline (Supplementary Fig. 6). The 25 ORFs₄ that are longer than 300 nucleotides, show signatures of translation and are under purifying selection, can confidently be considered genes despite being weakly conserved. The remaining 1,891 ORFs (1,139 ORFs₀⁺ and 752 ORFs₁₋₄) present characteristics intermediate between non-genic ORFs and genes, meeting our proto-gene designation (Fig. 4a, Supplementary Fig. 8 and Supplementary Table 3). We propose to place these ORFs in a continuum where strict annotation boundaries no longer have to be set (Fig. 4b).

Gene birth mechanisms involving re-organization of pre-existing genes, notably following gene duplication, have long been regarded as the predominant source of evolutionary innovation^{1,2}. Since the split between *S. cerevisiae* and *S. paradoxus*, sporadic gene duplications have generated between 1 and 5 novel genes²⁷. In contrast, 19 of the 143 ORFs₁ that arose *de novo* during the same evolutionary period were found under purifying selection. Therefore, *de novo* gene birth appears more prevalent than previously supposed^{3,10,12}, in agreement with recent estimations in humans and other primates^{1,9}. The involvement of proto-genes in *de novo* emergence of protein-coding genes in *S. cerevisiae* likely holds for other species and may extend to RNA genes and regulatory elements. Examination of translation program remodelling upon stress, in light of our evolutionary model, may further understanding of phenotypic diversity and plasticity of cellular systems^{7,28}.

Methods Summary

Detection of translation signatures

The mapping of ribosome footprint reads to ORFs does not necessarily indicate full-length, ORF-specific translation events^{6,25}. To model the number of ORFs₀⁺ expected if the detected presence of ribosomes on non-genic sequences was not related to the presence of ORFs₀, we randomized the positions of ORFs₀ while maintaining their length distribution and the observed positions of RNAseq and footprint reads. To model the number of ORFs₀⁺ expected if footprint reads observed outside of annotated ORFs were non-specific, we randomized the positions of footprint reads throughout non-genic sequences while maintaining the length distribution of footprint reads, the positions of RNAseq reads and the positions of ORFs₀. We optimized three parameters with regard to these two null models: i) the proportion of ORF length covered in RNAseq and footprint reads was fixed at 50% minimum; ii) the factor by which the number of footprint reads per nucleotide in the ORF should be higher than the number of footprint reads per nucleotide in surrounding up- and downstream windows was fixed at a minimum of 5; iii) the size of these windows was fixed at 300 nucleotides. Any two ORFs₀ that partially overlap on the same strand and show translation signatures in the same experimental conditions were both eliminated from the set of ORFs₀ considered to show translation signatures.

Significant purifying selection signatures

We estimated the number of synonymous mutations per synonymous site (dS) and the number of non-synonymous mutations per non-synonymous site (dN) for each ORF present

without disruptive mutations in eight *S. cerevisiae* strains. The likelihood of the dN/dS ratio for each ORF present without disruptive mutations in eight *S. cerevisiae* strains was determined under two distinct null models: assuming neutral evolution (the rates of synonymous and non-synonymous substitutions are equal) and not assuming neutral evolution. All ORFs with dN/dS < 1 and $P < 0.05$ (chi-square distribution of likelihoods with one degree of freedom) were considered to be subject to significant purifying selection.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank L. Duret, E. Levy, J. Vandenhoute, Q. Li, H. Yu, P. Braun, M. Dreze, C. Foo, M. Mann, N. Kulak, J. Cox, C. Maire and S. Jhavery-Schneider as well as members of the Center for Cancer Systems Biology (CCSB) in particular A. Dricot-Ziter, A. MacWilliams, F. Roth, Y. Jacob and D. Hill for discussions and proofreading. We apologize for the omission of relevant studies that we could not cite due to space limitations. A.R. was supported by a National Institute of Health Pioneer Award, a Career Award at the Scientific Interface from the Burroughs Wellcome Fund and the Howard Hughes Medical Institute (HHMI). I.W. is a HHMI fellow of the Damon Runyon Cancer Research Institute. G.A.B. was supported by American Cancer Society Postdoctoral fellowship 117945-PF-09-136-01-RMC. M.V. is a Chercheur Qualifié Honoraire from the Fonds de la Recherche Scientifique (FRS-FNRS, Wallonia-Brussels Federation, Belgium). This work was supported by the grant R01-HG006061 from the National Human Genome Research Institute awarded to M.V.

References

1. Tautz D, Domazet-Lošo T. The evolutionary origin of orphan genes. *Nat. Rev. Genet.* 2011; 12:692–702. [PubMed: 21878963]
2. Kaessmann H. Origins, evolution, and phenotypic impact of new genes. *Genome Res.* 2010; 20:1313–1326. [PubMed: 20651121]
3. Jacob F. Evolution and tinkering. *Science.* 1977; 196:1161–1166. [PubMed: 860134]
4. Siepel A. Darwinian alchemy: Human genes from noncoding DNA. *Genome Res.* 2009; 19:1693–1695. [PubMed: 19797681]
5. Khalturin K, Hemmrich G, Fraune S, Augustin R, Bosch TC. More than just orphans: are taxonomically-restricted genes important in evolution? *Trends Genet.* 2009; 25:404–413. [PubMed: 19716618]
6. Wilson BA, Masel J. Putatively noncoding transcripts show extensive association with ribosomes. *Genome Biol. Evol.* 2011; 3:1245–1252. [PubMed: 21948395]
7. Jarosz DF, Taipale M, Lindquist S. Protein homeostasis and the phenotypic manifestation of genetic diversity: principles and mechanisms. *Annu. Rev. Genet.* 2010; 44:189–216. [PubMed: 21047258]
8. Cai J, Zhao R, Jiang H, Wang W. *De novo* origination of a new protein-coding gene in *Saccharomyces cerevisiae*. *Genetics.* 2008; 179:487–496. [PubMed: 18493065]
9. Wu DD, Irwin DM, Zhang YP. *De novo* origin of human protein-coding genes. *PLoS Genet.* 2011; 7:e1002379. [PubMed: 22102831]
10. Ekman D, Elofsson A. Identifying and quantifying orphan protein sequences in fungi. *J. Mol. Biol.* 2010; 396:396–405. [PubMed: 19944701]
11. Lipman DJ, Souvorov A, Koonin EV, Panchenko AR, Tatusova TA. The relationship of protein conservation and sequence length. *BMC Evol. Biol.* 2002; 2:20. [PubMed: 12410938]
12. Wolf YI, Novichkov PS, Karev GP, Koonin EV, Lipman DJ. The universal distribution of evolutionary rates of genes and distinct characteristics of eukaryotic genes of different apparent ages. *Proc. Natl. Acad. Sci. USA.* 2009; 106:7273–7280. [PubMed: 19351897]
13. Cai JJ, Petrov DA. Relaxed purifying selection and possibly high rate of adaptation in primate lineage-specific genes. *Genome Biol. Evol.* 2010; 2:393–409. [PubMed: 20624743]

14. Zheng D, Gerstein MB. The ambiguous boundary between genes and pseudogenes: the dead rise up, or do they? *Trends Genet.* 2007; 23:219–224. [PubMed: 17382428]
15. Oliver SG, et al. The complete DNA sequence of yeast chromosome III. *Nature.* 1992; 357:38–46. [PubMed: 1574125]
16. Fisk DG, et al. *Saccharomyces cerevisiae* S288C genome annotation: a working hypothesis. *Yeast.* 2006; 23:857–865. [PubMed: 17001629]
17. Nagalakshmi U, et al. The transcriptional landscape of the yeast genome defined by RNA sequencing. *Science.* 2008; 320:1344–1349. [PubMed: 18451266]
18. Boyer J, et al. Large-scale exploration of growth inhibition caused by overexpression of genomic fragments in *Saccharomyces cerevisiae*. *Genome Biol.* 2004; 5:R72. [PubMed: 15345056]
19. Brar GA, et al. High-resolution view of the yeast meiotic program revealed by ribosome profiling. *Science.* 2012; 335:552–557. [PubMed: 22194413]
20. Li QR, et al. Revisiting the *Saccharomyces cerevisiae* predicted ORFeome. *Genome Res.* 2008; 18:1294–1303. [PubMed: 18502943]
21. Jansen R, Gerstein M. Analysis of the yeast transcriptome with structural and functional categories: characterizing highly expressed proteins. *Nucleic Acids Res.* 2000; 28:1481–1488. [PubMed: 10684945]
22. Giacomelli MG, Hancock AS, Masel J. The conversion of 3' UTRs into coding regions. *Mol. Biol. Evol.* 2007; 24:457–464. [PubMed: 17099057]
23. Prat Y, Fromer M, Linial N, Linial M. Codon usage is associated with the evolutionary age of genes in metazoan genomes. *BMC Evol. Biol.* 2009; 9:285. [PubMed: 19995431]
24. Yomtovian I, Teerakulkittipong N, Lee B, Moulton J, Unger R. Composition bias and the origin of ORFan genes. *Bioinformatics.* 2010; 26:996–999. [PubMed: 20231229]
25. Ingolia NT, Ghaemmaghami S, Newman JR, Weissman JS. Genome-wide analysis *in vivo* of translation with nucleotide resolution using ribosome profiling. *Science.* 2009; 324:218–223. [PubMed: 19213877]
26. Vishnoi A, Kryazhimskiy S, Bazykin GA, Hannenhalli S, Plotkin JB. Young proteins experience more variable selection pressures than old proteins. *Genome Res.* 2010; 20:1574–1581. [PubMed: 20921233]
27. Gao LZ, Innan H. Very low gene duplication rate in the yeast genome. *Science.* 2004; 306:1367–1370. [PubMed: 15550669]
28. Hayden EJ, Ferrada E, Wagner A. Cryptic genetic variation promotes rapid evolutionary adaptation in an RNA enzyme. *Nature.* 2011; 474:92–95. [PubMed: 21637259]
29. Pal C, Papp B, Hurst LD. Highly expressed genes in yeast evolve slowly. *Genetics.* 2001; 158:927–931. [PubMed: 11430355]
30. Drummond DA, Raval A, Wilke CO. A single determinant dominates the rate of yeast protein evolution. *Mol. Biol. Evol.* 2006; 23:327–337. [PubMed: 16237209]

Information)¹. Top right: number of ORFs assigned to each conservation level (logarithmic scale).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Fig. 2. Existence of an evolutionary continuum ranging from non-genic ORFs to genes through proto-genes

a, Length (top; error bars represent s.e.m.), RNA expression level (middle; error bars represent s.e.m.), and proximity to transcription factor binding sites (bottom; error bars represent standard error of the proportion) of ORFs correlate with conservation level. P and τ : Kendall's correlation statistics. Estimation of RNA abundance from RNAseq²⁵ in rich conditions. The positive correlation between proximity to transcription factor binding sites and conservation level is shown for a window of 200 nucleotides and holds when considering windows of 300, 400 and 500 nucleotides (Kendall's $\tau = 0.14, 0.16, 0.17$, respectively; $P < 2.2 \times 10^{-16}$ in each case). **b**, Codon bias increases with conservation level. Codon bias estimated using the codon adaptation index (Supplementary Information). P and τ : Kendall's correlation statistics. Error bars represent s.e.m. The large s.e.m. observed for ORFs₅ may be related to the whole genome duplication event (Supplementary Fig. 3). **c**, Relative amino acid abundances shift with increasing conservation level. For each encoded amino acid, the ratio between its frequency in ORFs₁₋₄ and its frequency in ORFs₅₋₁₀ (gray), or the ratio between its frequency in ORFs₁₋₄ and its frequency in ORFs₀ (black), is plotted. Enrichment of cysteine in proteins encoded by ORFs₁₋₄ relative to those encoded by ORFs₅₋₁₀ ($P < 1.8 \times 10^{-150}$, hypergeometric test) corresponds to 3.6 ± 0.1 residues (mean, s.e.m.) per translation product. **d**, Predicted structural features of ORF translation products correlate with conservation level. ORFs₀ were not included in these analyses as their short length hinders the reliability of structural predictions. Error bars represent s.e.m.




Fig. 3. Translation and adaptive potential of recently emerged ORFs

a, Example of an ORFs₀⁺ showing signatures of translation in starvation conditions. Syntenic regions in *Saccharomyces sensu stricto* species are aligned. Orange and black boxes: in-frame start and stop sites, respectively; *SCER*: *S. cerevisiae*; *SPAR*: *S. paradoxus*; *SMIK*: *S. mikatae*; *SBAY*: *S. bayanus*. **b**, Significance of the observed number of ORFs₀⁺. Distribution of the number of ORFs₀ expected to show signatures of translation if the ribosome footprinting assay were non specific (as modelled by randomizing footprint reads positions 100 times; squares), or if the presence of ribosomes on non-genic transcripts were not related to the presence of ORFs₀ (as modelled by randomizing ORFs₀ positions 100 times; circles). *P*: empirical *P* value. **c**, AUG context of ORFs with and without translation signatures. The presence of an adenine at position -3 from the start codon indicates optimum AUG context (Supplementary Information). *P* and *tau*: Kendall's correlation statistics. Asterisks (*) mark significant differences between ORFs with and without translation signatures ($P < 0.05$, Fisher's exact test). **d**, Candidate proto-genes tend to undergo condition-specific translation. **e**, Signatures of intra-species purifying selection. The positive correlation holds when only considering ORFs that are free from overlap with ORFs₁₋₁₀ (Supplementary Fig. 7), and is not entirely driven by the interdependence between strength of purifying selection and expression level (Supplementary Information)^{29,30}. Asterisk (*) marks a significant difference in proportion of ORFs under significant intra-species purifying selection between ORFs₀⁺ and ORFs₁ ($P = 0.0001$, hypergeometric test). *P* and *tau*: Kendall's correlation statistics. Error bars represent standard error of the proportion in all panels.




Fig. 4. Identification of proto-genes in a continuum ranging from non-genic ORFs to genes
a, Characterization of candidate proto-genes (ORFs₀⁺ and ORFs₁₋₄). Venn diagram not drawn to scale. **b**, The binary model of annotation (top) and the proposed continuum (bottom).